

Physics 121.
Lecture 15.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

1

Physics 121.
Lecture 15.

- Course Information
- Quiz.
- Topics to be discussed today:
 - Variables used to describe rotational motion
 - The equations of motion for rotational motion

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

2

Course Announcements.

- Homework set # 6 is now available on the WEB and will be due on Saturday morning, April 4, at 8.30 am.
- There will be no additional office hours and workshops this week on Tuesday, Wednesday, Thursday, and Friday.
- Exam # 2 will be returned to you during workshops next week.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

3

Quiz.

- Tell me about Exam # 2.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

4

Rotational motion: variables.

- In our discussion of rotational motion we will first focus on the rotation of **rigid** objects around a **fixed** axis.
- The variables that are used to describe this type of motion are similar to those we use to describe linear motion:
 - Angular position
 - Angular velocity
 - Angular acceleration




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

5

Rotational motion: variables.

- The angular position, measured in radians, is the angle of rotation of the object with respect to a reference position.
- The angular position of point P at this point in time is equal to θ . In order to uniquely define this position, we have assumed that the angular position is measured with respect to the x axis.



Frank L. H. Wolfs


Department of Physics and Astronomy, University of Rochester

6

Rotational motion: variables.

- Note:

- The angular position is always specified in radians!!!!
- One radian is the angular displacement corresponding to a linear displacement $l = R$.
- Make sure you keep track of the sign of the angular position!!!!
- An increase in the angular position corresponds to a counter-clockwise rotation; a decrease corresponds to a clockwise rotation.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

7

Complex motion in Cartesian coordinates is simple motion in rotational coordinates.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

8

Rotational motion: variables.

- If we look at an object carrying out a rotation around a fixed axis, we will see that the angular position becomes a function of time.
- To describe the rotational motion, we introduce the concepts of angular velocity and angular acceleration.
- **Remember:** for linear motion we found it useful to introduce the concepts of linear velocity and linear acceleration.



Angular acceleration (rad/s^2)

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester


9

Rotational motion: variables.

- For both velocity and acceleration, we can talk about instantaneous and average.

- Angular velocity:
 - Definition: $\omega = \frac{d\theta}{dt}$
 - Symbol: ω
 - Units: rad/s

- Angular acceleration:
 - Definition: $\alpha = \frac{d\omega}{dt} = \frac{d^2\theta}{dt^2}$
 - Symbol: α
 - Units: rad/s²



Angular acceleration (rad/s²)

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

10


Rotational motion: constant acceleration.

- If the object experiences a constant angular acceleration, then we can describe its rotational motion with the following equations of motion:

$$\omega(t) = \omega_0 + \alpha t$$

$$\theta(t) = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2$$

- Note how similar these equations are to the equation of motion for linear motion!!!!



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

11

Rotational motion: constant acceleration. Example problem.

- A wheel starting from rest, rotates with a constant angular acceleration of 2.0 rad/s². During a certain 3.0 s interval it turns through 90 rad. (a) How long had the wheel been turning before the start of the 3.0 s interval? (b). What was the angular velocity of the wheel at the start of the 3.0 s interval?
- Define time $t = 0$ s as the time that the wheel is at rest. The angular velocity and the angle of rotation at a later time t are given by

$$\omega(t) = \alpha t \quad \theta(t) = \frac{1}{2} \alpha t^2$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

12

Rotational motion: constant acceleration. Example problem.

- The change in the angular position $\Delta\theta$ during a time period Δt can now be calculated:

$$\Delta\theta(t) = \theta(t + \Delta t) - \theta(t) = \frac{1}{2}\alpha(\Delta t)^2 + \alpha t \Delta t$$

- Since the problem specifies $\Delta\theta$ and Δt we can now calculate the time t and the angular velocity at that time:

$$t = \frac{\Delta\theta - \frac{1}{2}\alpha(\Delta t)^2}{\alpha\Delta t} \quad \omega = \alpha t = \frac{\Delta\theta - \frac{1}{2}\alpha(\Delta t)^2}{\Delta t} = \frac{\Delta\theta}{\Delta t} - \frac{1}{2}\alpha\Delta t$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

13



5 Minute 16 Second Intermission.

- Since paying attention for 1 hour and 15 minutes is hard when the topic is physics, let's take a 5 minute 16 second intermission.

- You can:
 - Stretch out.
 - Talk to your neighbors.
 - Ask me a quick question.
 - Enjoy the fantastic music.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

14

Rotational motion: variables.

- The linear velocity of a part of the rigid body is related to the angular velocity of the object.


- Consider point P:

- It this point makes one complete revolution, it travels a distance $2\pi R$.
- When the angular position changes by $\Delta\theta$, point P moves a distance

$$\Delta l = 2\pi R \frac{\Delta\theta}{2\pi} = R\Delta\theta$$

- The linear velocity of point P is equal to

$$v = \frac{dl}{dt} = R \frac{d\theta}{dt} = R\omega$$



Frank L. H. Wolfs


Department of Physics and Astronomy, University of Rochester

15

Rotational motion: variables.

- Things to consider when looking at the rotation of rigid objects around a fixed axis:

- Each part of the rigid object has the same angular velocity.
- Only those parts that are located at the same distance from the rotation axis have the same linear velocity.
- The linear velocity of parts of the rigid object increases with increasing distance from the rotation axis.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

16


Relation between rotational and linear variables.

- Although in rotational motion we prefer to use rotational variables, we can also express the motion in terms of linear variables:

$$s = r\theta$$

$$v = \frac{ds}{dt} = \frac{d}{dt}(r\theta) = r \frac{d\theta}{dt} = r\omega$$

$$a_t = \frac{dv}{dt} = \frac{d}{dt}(r\omega) = r \frac{d\omega}{dt} = r\alpha$$



Frank L. H. Wolfs


Department of Physics and Astronomy, University of Rochester

17

Rotational motion: acceleration.

- Note: the acceleration $a_t = r\alpha$ is only one of the two components of the acceleration of point P. The two components of the acceleration of point P are:

- The radial component: this component is always present since point P carried out circular motion around the axis of rotation.
- The tangential component: this component is present only when the angular acceleration is not equal to 0 rad/s².




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

18

Rolling motion.

- To describe rolling motion we need to use both translational and rotational motion.
- The rolling motion can be described in terms of pure rotational motion with respect to the contact point P which is always at rest.
- The rotation axis around P is called the instantaneous axis and will move when the wheel rolls.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

19

Direction of the angular velocity. Use your right hand!

Angular velocity and acceleration are vectors!
They have a magnitude and a direction.



Frank L. H. Wolfs


Department of Physics and Astronomy, University of Rochester

20

Direction of the angular acceleration.

The angular acceleration α is parallel or anti-parallel to the angular velocity:

- If ω increases: parallel
- If ω decreases: anti-parallel




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

21

Pseudo vectors.

Angular velocity and accelerations are not real vectors!
They do not behave as real vectors under reflection: they are what we call pseudo vectors.




Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

22

Done for today!
Much more about rotations on Tuesday!



Opportunity rover indicates ancient Mars was wet.
Credit: [Mars Exploration Rover Mission](#), [JPL](#), [NASA](#).

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester

23
